

Directions: 1. Indicate choice when options are available by placing a check in the box
 2. Mark through entire line any prechecked item you ***do not*** wish to order ~~mark through~~

Attending Physician: _____ Diagnosis: _____
 Observation Status Admission for Inpatient Care Admit or transfer to: _____ Unit

ALLERGIES: NKA **ALLERGIC to:** _____

VITAL SIGNS:

- Vital signs on admission
- Vital signs every 1 hour pre-operatively
- Vital signs every 30 minutes if pregnancy induced hypertension
- Vital signs every 15 minutes x 4 post-operatively; THEN
- Vital signs every 30 minutes post-operatively

DIET:

- NPO

IV SOLUTIONS:

- Lactated Ringers IV 1000 ml bolus infuse over 1 hour; THEN
- Lactated Ringers IV infuse at 150 ml/hour
- Lactated Ringers IV 1000 ml with **oxytocin (Pitocin)** 20 units bolus infuse over 1 hour post-operatively

MEDICATIONS:

- Cefazolin (Ancef)** 2 grams IVPB x 1 dose, for wound infection prophylaxis
- Clindamycin** 900 mg IVPB x 1 dose, for wound infection prophylaxis

Pain medication for patient with **SPINAL/EPIDURAL ANESTHESIA**

- Morphine** 2.5 mg IV x 1 dose may repeat in 20 mins PRN for pain; THEN
- Morphine** 2.5 mg IV every 3 hours x 12 hours PRN for pain; THEN
- Morphine** 5 mg IV every 3 hours PRN for pain

Pain medication for patient with **GENERAL ANESTHESIA**

- Morphine** 5 mg IV every 3 hours PRN for pain
- Ondansetron (Zofran)** 4 mg IV every 4 hours PRN for nausea
- Diphenhydramine (Benadryl)** 25 mg IV every 2 hours PRN x 2 doses for severe pruritus post delivery

PATIENT CARE:

- External fetal monitoring
- Indwelling catheter to straight bedside drainage
- Abdominal prep
- Sequential compression devices (SCD) place prior to surgery

LABORATORY:

- CBC with Differential (stat)
- RPR (rapid plasma reagin) if status unknown
- PT/INR with admission labs (if patient has been on anticoagulant therapy)
- aPTT with admission labs (if patient has been on anticoagulant therapy)
- PFA (platelet function assay) with admission labs (if patient has been on anticoagulant therapy)

COMMUNICATION:

- Obtain documentation of GBS (Group B Streptococcus) status
- Obtain cord blood and send to lab if patient Rh negative or O+
- Recovery period to be 1-2 hours after delivery, may be extended if indicated

ADDITIONAL ORDERS:

Physician _____	Pager _____	Date ____ / ____ / ____	Time: _____
<input type="checkbox"/> Verbal Order Verification Signature _____		Date ____ / ____ / ____	Time: _____