

Directions: 1. Indicate choice when options are available by placing a check in the box
 2. Mark through entire line any prechecked item you ***do not*** wish to order ~~mark through~~

Attending Physician: _____ Diagnosis: _____
 Observation Status Admission for Inpatient Care Admit or transfer to: _____ Unit

ALLERGIES: NKA **ALLERGIC to:** _____

VITAL SIGNS:

- Vital signs every 1 hour
- Vital signs every 30 minutes if Pregnancy Induced Hypertension (PIH)
- Vital signs every 15 minutes x 4 post delivery

ACTIVITY:

- Ambulate as tolerated during early labor
- Bed rest

DIET:

- NPO, except ice chips

IV SOLUTIONS:

- Lactated Ringers IV infuse 1 liter over 1 hour prior to regional anesthesia
- Lactated Ringers IV infuse at 150 ml/hour
- Dextrose 5% and Lactated Ringers IV infuse at 150 ml/hour
- Pitocin Induction/Augmentation: Lactated Ringers 250 ml and **pitocin (Oxytocin)** 15 units IV infuse per low dose titration protocol
- Lactated Ringers 1000 ml with **pitocin (Oxytocin)** 20 units bolus infuse over 1 hour post delivery
- Change IV to saline lock prior to transfer post delivery

MEDICATIONS:

- Cervical Induction: Insert **dinoprostone (Cervidil)** 10 mg intravaginally in posterior fornix of cervix
- Ampicillin** 2 grams IVPB every 6 hours; indication: GBS prophylaxis
- Ampicillin** 2 grams IVPB to load (indication: GBS Prophylaxis); THEN
- Ampicillin** 1gram IVPB every 4 hours; indication: GBS prophylaxis
- Cefazolin(Ancef)** 2 grams IVPB every 8 hours; indication: GBS prophylaxis
- Clindamycin** 900 mg IVPB every 8 hours; indication: GBS prophylaxis
- Penicillin** 5 million units IVPB to load (indication: GBS prophylaxis); THEN
- Penicillin** 2.5 million units IVPB every 4 hours; indication: GBS prophylaxis
- Butorphanol tartrate (Stadol)** 1 mg IV every 1 hour PRN for pain level of mild to moderate
NOT TO EXCEED 24 MG OF **STADOL** IN 24 HOURS
- Morphine** 5 mg IV push every 3 hours PRN for pain level of severe

Directions: 1. Indicate choice when options are available by placing a check in the box
 2. Mark through entire line any prechecked item you ***do not*** wish to order ~~mark through~~

MEDICATIONS: Cont'd

- Ondansetron (Zofran)** 4 mg IV every 4 hours PRN for nausea
- Hydroxyzine pamoate (Vistaril)** 100 mg PO PRN at bedtime for rest
- Zolpidem (Ambien)** 10 mg PO PRN at bedtime for rest
- Oxycodone/acetaminophen (Percocet)** 5/325 mg one (1) tablet PO every 4 hours PRN for mild pain post delivery
- Oxycodone/acetaminophen (Percocet)** 5/325, two (2) tablets PO every 4 hours PRN for moderate to severe pain post delivery
- Ibuprofen (Motrin)** 800 mg PO every 8 hours PRN for mild to severe pain post delivery

PATIENT CARE:

- External fetal monitoring
- Straight urinary catheter PRN for bladder distention, if unable to void
- Ice pack PRN to perineum for patient comfort post delivery
- Fundus height and bleeding check every 15 minutes x 4, then per unit protocol post delivery

LABORATORY:

- CBC with Differential stat
- Type and screen
- RPR (rapid plasma reagin) if status unknown
- PT/INR with admission labs (if patient has been on anticoagulant therapy)
- aPTT with admission labs (if patient has been on anticoagulant therapy)
- PFA (platelet function assay) with admission labs (if patient has been on antiplatelet therapy)
- Rubella Antibody IgG (if status unknown)

COMMUNICATION:

- Obtain documentation of GBS (Group B Streptococcus) status
- Obtain cord blood and send to lab on miscellaneous paper order If patient Rh negative or O+

ADDITIONAL ORDERS:

Physician _____	Pager _____	Date ____ / ____ / ____	Time: _____
<input type="checkbox"/> Verbal Order Verification Signature _____		Date ____ / ____ / ____	Time: _____

****SCAN PHYSICIAN SIGNATURE PAGE FIRST TO PHARMACY****